

Gene Section

Mini Review

NUP98 (nucleoporin 98 kDa)

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France (JLH)

Published in Atlas Database: November 1998

Online updated version : <http://AtlasGeneticsOncology.org/Genes/NUP98.html>
DOI: 10.4267/2042/37478

This article is an update of: Huret JL. NUP98 (nucleoporin 98 kDa). *Atlas Genet Cytogenet Oncol Haematol.* 1998;2(1):7.

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 1999 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

Identity

Location : 11p15

DNA/RNA

Transcription

3.6, 6.5, 7.0 kb mRNA.

Protein

Description

920 amino acids; 97 kDa; contains repeated motifs (GLFG and FG) in N-term and a RNA binding motif in C-term.

Expression

Wide.

Localisation

Nuclear membrane localisation.

Function

Nucleoporin: associated with the nuclear pore complex; role in nucleocytoplasmic transport processes.

Homology

Member of the GLFG nucleoporins.

Implicated in

t(7;11)(p15;p15)/ANLL --> NUP98/HOXA9

Disease

M2-M4 ANLL mostly; occasionally, CML-like cases.

Prognosis

Mean survival: 15 months.

Cytogenetics

Sole anomaly most often.

Hybrid/Mutated gene

5' NUP98 - 3' HOXA9.

Abnormal protein

Fuses the GLFG repeat domains of NUP98 to the HOXA9 homeobox.

inv(11)(p15q22)/MDS or ANLL --> NUP98/DDX10

Disease

Therapy-related MDS and ANLL; de novo ANLL.

Hybrid/Mutated gene

5' NUP98 - 3' DDX10

Abnormal protein

Fuses the GLFG repeat domains of NUP98 to the acidic domain of DDX10.

t(2;11)(q31;p15)/treatment-related leukaemia --> NUP98/HOXD13

Disease

So far, only 1 case of treatment related myelodysplasia evolving towards M6 acute non lymphocytic leukaemia.

Hybrid/Mutated gene

5' NUP98 - 3' HOXD13.

Abnormal protein

Fuses the GLFG repeat domains of NUP98 to the HOXD13 homeodomain.

References

Nakamura T, Largaespada DA, Lee MP, Johnson LA, Ohyashiki K, Toyama K, Chen SJ, Willman CL, Chen IM, Feinberg AP, Jenkins NA, Copeland NG, Shaughnessy JD Jr. Fusion of the nucleoporin gene NUP98 to HOXA9 by the chromosome translocation t(7;11)(p15;p15) in human myeloid leukaemia. *Nat Genet.* 1996 Feb;12(2):154-8

Arai Y, Hosoda F, Kobayashi H, Arai K, Hayashi Y, Kamada N, Kaneko Y, Ohki M. The inv(11)(p15q22) chromosome translocation of de novo and therapy-related myeloid malignancies results in fusion of the nucleoporin gene, NUP98, with the putative RNA helicase gene, DDX10. *Blood.* 1997 Jun 1;89(11):3936-44

Powers MA, Forbes DJ, Dahlberg JE, Lund E. The vertebrate GLFG nucleoporin, Nup98, is an essential component of multiple RNA export pathways. *J Cell Biol.* 1997 Jan 27;136(2):241-50

Raza-Egilmez SZ, Jani-Sait SN, Grossi M, Higgins MJ, Shows TB, Aplan PD. NUP98-HOXD13 gene fusion in therapy-related acute myelogenous leukemia. *Cancer Res.* 1998 Oct 1;58(19):4269-73

This article should be referenced as such:

Huret JL. NUP98 (nucleoporin 98 kDa). *Atlas Genet Cytogenet Oncol Haematol.* 1999; 3(1):15-16.