

CARACTÉRISATION PHYTOSOCIOLOGIQUE DES CHÊNAIES LITTORALES DU FINISTÈRE

Frédéric BIORET - Sébastien GALLET

Tableau A *Rubio peregrinae-Quercetum petraeae* ass. nov. hoc loco

| N° des relevés | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | Σ |
|-------------------------------|---------------|--------------|---------------|---------------|-------------|--------------|---------------|----------------|-------------|-------------|---------------|---------------|-----|
| Surface (m2) | 400 | 500 | 400 | 400 | 300 | 400 | 500 | 600 | 500 | 400 | 500 | 500 | |
| Exposition | N NW | W | SW | S-SW | S-SW | SW | N | N | N-NE | N-NE | E-NE | N-NE | |
| A | 10-12m 90% | 8-12m 95% | 12-14m 80% | 10-15m 95% | 6-8m, 90% | 8-10m 85% | 10-12m 90% | 10-15m, 80% | 12-14m, 80% | 10-15m, 80% | 15m, 80% | 12-15m 80% | |
| <i>Quercus petraea</i> | 55 | 55 | 54 | 55 | 55 | 55 | 55 | 55 | 54 | 54 | 55 | 55 | V |
| <i>Quercus robur</i> | + | | 11 | + | | | | | | + | | | II |
| <i>Castanea sativa</i> | | | | | | | | i | | | | | I |
| <i>Hedera helix</i> | | | | | | | 11 | + | + | | | + | |
| a | 3-6m, 40% | 3-6m, 50% | 2-4m, 30% | 2-3m, 40% | 2-4m, 15% | 3-4m, 20% | 4-6m, 30% | 3-6m, 40% | 3-4m, 15% | 3-4m, 25% | 2-5m, 20% | 1,5-3m, 5% | |
| <i>Ilex aquifolium</i> | 34 | 44 | 33 | 33 | +2 | 23 | 22 | 33 | + | 22 | 22 | + | V |
| <i>Sorbus aucuparia</i> | | | | | | | | 11 | 11 | 11 | 11 | 12 | III |
| <i>Sorbus torminalis</i> | 23 | + | | | | | 22 | + | | | | | II |
| <i>Mespilus germanica</i> | + | 11 | | + | + | | | | | | | | II |
| <i>Corylus avellana</i> | | | | | | | | 22 | 11 | | +2 | | II |
| <i>Pyrus cordata</i> | | | +2 | | + | + | | | | | | + | II |
| <i>Quercus petraea</i> | + | 11 | 12 | | | | | | | | | 11 | II |
| <i>Hedera helix</i> | | | 12 | | | | | | + | + | + | | II |
| <i>Ulex europaeus</i> | | | | | i | | | | | | | + | I |
| <i>Tilia cordata</i> | | | | | | | | | + | | | | + |
| <i>Rhamnus frangula</i> | | | | | | | | | | | + | | + |
| <i>Fagus sylvatica</i> | | | | | | | i | | | | | | + |
| <i>Prunus avium</i> | | | | | + | | | | | | | | + |
| <i>Lonicera periclymenum</i> | | | | | 11 | | | | | | | | + |
| h | 0,5-1m, 75% | 0,1-1m, 60% | 0,1-1m, 90% | 0,1-1m, 90% | 0,1-1m, 60% | 0,1-1m, 70% | 0,2-1m, 100% | 0,1-0,9m, 100% | 0,1-1m, 80% | 0,2-1m, 90% | 0,15-1m, 100% | 0,3-1,1m, 80% | |
| <i>Vaccinium myrtillus</i> | +2 | + | | | | 33 | + | +2 | 22 | 11 | 32 | 22 | IV |
| <i>Luzula sylvatica</i> | | | | | | | 55 | 55 | 55 | 54 | 55 | 44 | III |
| <i>Lonicera periclymenum</i> | + | 11 | 22 | + | + | 22 | 11 | + | 11 | 11 | | 11 | V |
| <i>Rubus sp.</i> | 33 | + | 22 | +2 | | | 22 | 22 | 12 | 22 | 11 | + | V |
| <i>Hedera helix</i> | 33 | 22 | 33 | | | | 11 | + | 11 | + | 11 | 11 | IV |
| <i>Ruscus aculeatus</i> | 23 | +2 | 44 | 12 | 22 | +2 | 11 | +2 | + | | | | IV |
| <i>Thuidium tamariscinum</i> | 33 | 34 | 22 | 22 | 11 | + | + | | + | | | | IV |
| <i>Pteridium aquilinum</i> | + | + | | | | 11 | 11 | 11 | 22 | | 22 | + | IV |
| <i>Teucrium scorodonia</i> | + | | 23 | + | | | + | 11 | | + | | + | III |
| <i>Quercus petraea</i> | 11 | | + | 11 | | + | 11 | | | | + | +j | III |
| <i>Rubia peregrina</i> | 11 | + | + | + | 22 | | | | + | | | | III |
| <i>Melampyrum pratense</i> | | | | | + | 22 | | | + | 11 | | 33 | III |
| <i>Calluna vulgaris</i> | | + | | + | + | 12 | | | | | | + | II |
| <i>Polypodium interjectum</i> | | | | | | | + | | + | + | | + | II |
| <i>Polytrichum formosum</i> | | | +2 | + | + | | | | | | + | | II |
| <i>Sorbus aucuparia</i> | | | | | | + | | +j | | | + | +j | II |
| <i>Ilex aquifolium</i> | + | | | + | + | + | | | | | | | II |
| <i>Carex sp.</i> | + | | + | + | | | | | | | | | II |

| | | | | | | | | | |
|----------------------------------|---|----|---|----|----|----|----|---|----|
| <i>Dicranum scoparium</i> | | 11 | | | + | | + | | II |
| <i>Leucobryum glaucum</i> | | | | 12 | +2 | 11 | | | II |
| <i>Hyacinthoides non-scripta</i> | | | + | + | | | | | I |
| <i>Erica cinerea</i> | | + | | | | + | | | I |
| <i>Rhytiadelphus squarrosus</i> | | 11 | | | | | + | | I |
| <i>Fraxinus excelsior</i> | + | lj | | | | | | | I |
| <i>Corylus avellana</i> | + | | | | | | | + | I |
| <i>Fagus sylvatica</i> | i | | | | +j | | | | I |
| <i>Dryopteris filix-mas</i> | + | | | | | | | | + |
| <i>Ulex europaeus</i> | | + | | | | | | | + |
| <i>Cytisus scoparius</i> | | | + | | | | | | + |
| <i>Castanea sativa</i> | | | | | +j | | | | + |
| <i>Umbilicus rupestris</i> | | | | | | + | | | + |
| <i>Dactylis glomerata</i> | | | | | | + | | | + |
| <i>Anthoxanthum odoratum</i> | | | | | | + | | | + |
| <i>Brachypodium sylvaticum</i> | | | | | | | 11 | | + |
| <i>Plagothecium undulatum</i> | | | | | | | | + | + |

Localisation des relevés :

1, 2 , 7, 8 : Port Hascoët, Hôpital-Camfrout

3, 4, 5, 6 : Térénez, rive droite de l'Aulne

9, 10, 11, 12 : Forêt de Landévennec, rive gauche de l'Aulne