

Leukaemia Section

Short Communication

t(14;14)(q11;q32) CEBPE/IGH, inv(14)(q11q32) CEBPE/IGH

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France (JLH)

Published in Atlas Database: May 2008

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/t1414q11q32CEBPEIGHID1438.html>
DOI: 10.4267/2042/44476

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 2009 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Identity

Note

This chromosome anomaly should not be confused with the t(14;14)(q11;q32)/inv(14)(q11q32) found in T-cell diseases, which implicates TCR alpha or TCR delta (14q11) and TCL1A (14q32).

Clinics and pathology

Disease

CD10+ acute lymphoblastic leukaemia (B-ALL).

Epidemiology

Only 4 cases to date of t(14;14)(q11;q32)/inv(14)(q11q32) with CEBPE and IGH involvements (Akasaka et al., 2007). Five other cases of t(14;14)(q11;q32)/inv(14)(q11q32) in B-cell leukemias are known (Denny et al., 1986; Speleman et al., 1991; Chervinsky et al., 1995; Wong et al., 1995; Thomas et al., 2001), but without proof that CEBPE was involved. As a matter of fact, a t(4;11)(q21;q23) was found in 2 of these cases, and a t(8;14)(q24;q32) in another case; this latter group is certainly heterogeneous.

Clinics

The four patients were male patients, aged 15, 25, 45, and 45 years, with a WBC under $50 \times 10^9/l$. Survival is available only for two cases: 19 mths+ and 48 months+, resembling the relatively fair survival of

patients with a t(8;14)(q11;q32) CEBPD/IGH translocation.

One case was a Down syndrome patient; this may not be anecdotal, since more than 1/4 of t(8;14)(q11;q32) case are also Down syndrome patients.

Genes involved and proteins

CEBPE

Location

14q11

Protein

DNA-binding protein. CCAAT enhancer-binding protein (CEBP) transcription factors are a family of 6 multifunctional basic leucine zipper (bZIP) transcription factors. The 5 other CEBPs are: CEBPA (19q13), CEBPB (20q13), CEBPD (8q11), CEBPG (19q13), all four equally implicated in leukemias, and DDIT3/CHOP/CEBP zeta (12q13), so far known to be involved in solid tumours (liposarcoma). These transcription factors play a key role in cellular differentiation, in particular in the control of myeloid differentiation. CEBPE is composed of an N-term transactivation domain, a negative regulatory domain, a DNA-binding basic motif, and a leucine-zipper domain in C-term (Ramji et al., 2002; Nerlov et al., 2007).

IgH

Location

14q32

Result of the chromosomal anomaly

Fusion protein

Oncogenesis

Overexpression of the CEBP gene.

References

Denny CT, Hollis GF, Hecht F, Morgan R, Link MP, Smith SD, Kirsch IR. Common mechanism of chromosome inversion in B- and T-cell tumors: relevance to lymphoid development. *Science*. 1986 Oct 10;234(4773):197-200

Speleman F, Mangelschots K, Vercruyssen M, Dal Cin P, Aventin A, Offner F, Laureys G, Van den Berghe H, Leroy J. Analysis of whole-arm translocations in malignant blood cells by nonisotopic in situ hybridization. *Cytogenet Cell Genet*. 1991;56(1):14-7

Chervinsky DS, Grossi M, Kakati S, Block AW, Aplan PD. Concurrent presence of inv(14)(q11q32) and t(4;11)(q21;q23) in pre-B acute lymphoblastic leukemia. *Genes Chromosomes Cancer*. 1995 Mar;12(3):229-36

Wong KF, Kwong YL, Wong TK. Inversion 14q in acute lymphoblastic leukemia of B-lineage. *Cancer Genet Cytogenet*. 1995 Mar;80(1):72-4

Thomas X, Olteanu N, Charrin C, Lhéritier V, Magaud JP, Fiere D. Acute lymphoblastic leukemia in the elderly: The Edouard Herriot Hospital experience. *Am J Hematol*. 2001 Jun;67(2):73-83

Ramji DP, Foka P. CCAAT/enhancer-binding proteins: structure, function and regulation. *Biochem J*. 2002 Aug 1;365(Pt 3):561-75

Akasaka T, Balasas T, Russell LJ, Sugimoto KJ, Majid A, Walewska R, Karran EL, Brown DG, Cain K, Harder L, Gesk S, Martin-Subero JI, Atherton MG, Brüggemann M, Calasanz MJ, Davies T, Haas OA, Hagemeijer A, Kempski H, Lessard M, Lillington DM, Moore S, Nguyen-Khac F, Radford-Weiss I, Schoch C, Struski S, Talley P, Welham MJ, Worley H, Strefford JC, Harrison CJ, Siebert R, Dyer MJ. Five members of the CEBP transcription factor family are targeted by recurrent IGH translocations in B-cell precursor acute lymphoblastic leukemia (BCP-ALL). *Blood*. 2007 Apr 15;109(8):3451-61

Nerlov C. The C/EBP family of transcription factors: a paradigm for interaction between gene expression and proliferation control. *Trends Cell Biol*. 2007 Jul;17(7):318-24

This article should be referenced as such:

Huret JL. t(14;14)(q11;q32) CEBPE/IGH, inv(14)(q11q32) CEBPE/IGH. *Atlas Genet Cytogenet Oncol Haematol*. 2009; 13(5):376-377.
