

Gene Section

Mini Review

TAC1 (tachykinin, precursor 1)

Philip K Lim, Shyam A Patel, Pranela Rameshwar

University of Medicine and Dentistry of New Jersey - New Jersey Medical School, Newark, New Jersey, USA (PKL, SAP, PR)

Published in Atlas Database: June 2009

Online updated version : <http://AtlasGeneticsOncology.org/Genes/TAC1ID44483ch7q21.html>

DOI: 10.4267/2042/44761

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 2010 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Identity

Other names: NK2; NKA; NKNA; NPK; PPT; TAC2

HGNC (Hugo): TAC1

Location: 7q21.3

DNA/RNA

Description

The TAC1 gene maps (Homo sapiens) to NC_000007.12 in the region between 97199311 and 97207720 on the plus strand and spans 8410 bp.

Transcription

4 transcript variants:

alpha (1134 bp); open reading frame from bp 247-582
-lacks exon 6

-encodes substance P on exon 3

beta (1188 bp); open reading frame from bp 247-636

-encodes the full-length version of Tac1

-encodes substance P, neurokinin A, neuropeptide K

delta (1089 bp); open reading frame from bp 247-537

-lacks exons 4 and 6

-encodes substance P on exon 3

gamma (1143 bp); open reading frame from bp 247-591

-lacks exon 4

-encodes substance P, neurokinin A

Protein

Note

The TAC1 gene encodes multiple transcripts through post-translational modifications. Each encodes peptides belonging to the tachykinin family of peptides. The major peptides produced from the TAC1 transcripts are substance P and neurokinin A. Others include

neuropeptide K, and neuropeptide gamma. The tachykinins exert multiples functions such as neurotransmission, immune modulation and hematopoietic regulation. TAC1 encodes peptides that target nerve receptors, immune cells, stem cells, hematopoietic cells and smooth muscle cells. They function in vasodilatory responses; act as secretagogues and can induce behavioral responses.

Description

4 peptides:

alpha: 111 aa

-comprises substance P

beta: 129 aa

-comprises substance P, neurokinin A, neuropeptide K

delta: 96 aa

-comprises substance P

gamma: 114 aa

-comprises neuropeptide gamma, substance P, neurokinin A.

Expression

Expressed by various immune and neuronal cells.

TAC1 expression in the setting of osteoarthritis can be induced by mechanical stimulation (Howard et al., 2008).

TAC1 expression is also regulated by microRNAs in neurons derived from human mesenchymal stem cells (Greco et al., 2007). This occurs by binding of miRNA-13-a, miRNA206 and miRNA302a to the Tac1 3'UTR. HIV1 infection resulted in increase production of TAC1 peptide (substance P). The source of substance P has been identified as monocytes-derived macrophages from placenta cord blood and adult peripheral blood. The production of substance P correlates with HIV 1 infection (Douglas et al., 2002).

Localisation

Secreted peptide.

Function

TAC1 peptides function in both pathologic and physiologic processes. These include hematopoiesis, gastrointestinal secretory processes, respiratory patterns, calcium signaling, neuropeptide signaling, pain, synaptic transmission, insemination (Murthy et al., 2008; David et al., 2009); inflammatory response, autism, and pulmonary infection (Marui et al., 2007; Grissell et al., 2007). In normal respiratory development, TAC1 appears to be a crucial gene, exerting plasticity during development (Berner et al., 2007).

TAC1 is involved in hematopoietic regulation. The mRNA is targeted by RNA-binding protein that confers translational control, which could be negatively regulated by cytokines with hematopoietic stimulator properties (Murthy et al., 2008). TAC1 has been implicated in macrophage and monocyte functions (Chernova et al., 2009).

Homology

Homo sapiens TAC1 shares sequence homology with mouse and rat sequences.

Implicated in

Breast cancer

Note

A gene expression signature for breast cancer has been shown to involve TAC1 (Ellsworth et al., 2009).

Prognosis

TAC1 expression occurs in breast cancer and is directly proportional to aggressiveness of the cancer and thus, TAC1 may also be a prognostic factor in breast cancer (Ellsworth et al., 2009; Reddy et al., 2009).

Oncogenesis

TAC1 expression favors breast cancer cell entry into the bone marrow during stage IV disease (Reddy et al., 2009). TAC1 regulates the interaction between CXCL12 and its receptor, CXCR4 in the interaction between breast cancer and mesenchymal stem cells (Corcoran et al., 2008).

The TAC1 gene products, substance P and neurokinin A, and neurokinin receptor antagonists have been shown to have anti-proliferative effects on breast cancer cells (Rameswhar et al., 1996; Singh et al., 2000).

Colon cancer

Note

Substance P, the major peptide encoded by the TAC1 gene, is mitogenic to colon cancer. Its action on the cancer cells appears to be autocrine since NK1 antagonist has been shown to mediate anti-tumor activity (Rosso et al., 2008). On the other hand, Substance P also enhances the expansion of lymphokine-activated killer cells against colon cancer cells (Flageole et al., 1992). These two properties of

substance P appear paradoxical. Thus, targeted therapy will need to balance the immune-enhancing effects with the tumor promoting functions of substance.

Prognosis

In Dukes stage A/B cancers, TAC1 methylation levels were significantly higher than in Dukes stage C/D cancers (Mori et al., 2006).

Oncogenesis

TAC1 was found to be silenced via promoter methylation in primary colon cancer and may lead to early stage carcinogenesis by aiding tumor cells to escape immune surveillance and autocrine growth-inhibitory signaling (Mori et al., 2006).

Esophageal cancer

Prognosis

Promoter hypermethylation of TAC1 confers poor prognosis in esophageal cancer (Jin et al., 2007).

Oncogenesis

TAC1 promoter hypermethylation has been suggested to be a putative marker for esophageal carcinoma (Jin et al., 2007).

Gastric adenocarcinoma

Oncogenesis

Promoter hypermethylation of TAC1 has also been found in gastric adenocarcinoma (David et al., 2009).

Multiple sclerosis

Note

The genetic region surrounding TAC1 may be unstable and has been associated with increased susceptibility to multiple sclerosis (Vandenbroeck et al., 2002).

Narcolepsy

Note

TAC1 and other related peptides have been associated with narcolepsy. Stimulation by amphetamines promotes increased TAC1 expression (Lindberg et al., 2007).

References

- Flageole H, Senterman M, Trudel JL. Substance P increases in vitro lymphokine-activated-killer (LAK) cell cytotoxicity against fresh colorectal cancer cells. *J Surg Res.* 1992 Nov;53(5):445-9
- Rameshwar P, Gascón P. Induction of negative hematopoietic regulators by neurokinin-A in bone marrow stroma. *Blood.* 1996 Jul 1;88(1):98-106
- Singh D, Joshi DD, Hameed M, Qian J, Gascón P, Maloof PB, Mosenthal A, Rameshwar P. Increased expression of preprotachykinin-I and neurokinin receptors in human breast cancer cells: implications for bone marrow metastasis. *Proc Natl Acad Sci U S A.* 2000 Jan 4;97(1):388-93
- Ho WZ, Lai JP, Li Y, Douglas SD. HIV enhances substance P expression in human immune cells. *FASEB J.* 2002 Apr;16(6):616-8
- Vandenbroeck K, Fiten P, Heggarty S, Goris A, Cocco E, Hawkins SA, Graham CA, Marrosu MG, Opdenakker G.

Chromosome 7q21-22 and multiple sclerosis: evidence for a genetic susceptibility effect in vicinity to the protachykinin-1 gene. *J Neuroimmunol.* 2002 Apr;125(1-2):141-8

Mori Y, Cai K, Cheng Y, Wang S, Paun B, Hamilton JP, Jin Z, Sato F, Berki AT, Kan T, Ito T, Mantzur C, Abraham JM, Meltzer SJ. A genome-wide search identifies epigenetic silencing of somatostatin, tachykinin-1, and 5 other genes in colon cancer. *Gastroenterology.* 2006 Sep;131(3):797-808

Berner J, Shvarev Y, Lagercrantz H, Bilkei-Gorzo A, Hökfelt T, Wickström R. Altered respiratory pattern and hypoxic response in transgenic newborn mice lacking the tachykinin-1 gene. *J Appl Physiol.* 2007 Aug;103(2):552-9

Greco SJ, Rameshwar P. MicroRNAs regulate synthesis of the neurotransmitter substance P in human mesenchymal stem cell-derived neuronal cells. *Proc Natl Acad Sci U S A.* 2007 Sep 25;104(39):15484-9

Grissell TV, Chang AB, Gibson PG. Reduced toll-like receptor 4 and substance P gene expression is associated with airway bacterial colonization in children. *Pediatr Pulmonol.* 2007 Apr;42(4):380-5

Jin Z, Olaru A, Yang J, Sato F, Cheng Y, Kan T, Mori Y, Mantzur C, Paun B, Hamilton JP, Ito T, Wang S, David S, Agarwal R, Beer DG, Abraham JM, Meltzer SJ. Hypermethylation of tachykinin-1 is a potential biomarker in human esophageal cancer. *Clin Cancer Res.* 2007 Nov 1;13(21):6293-300

Lindberg J, Saetre P, Nishino S, Mignot E, Jazin E. Reduced expression of TAC1, PENK and SOCS2 in Hcrtr-2 mutated narcoleptic dog brain. *BMC Neurosci.* 2007 May 23;8:34

Marui T, Funatogawa I, Koishi S, Yamamoto K, Matsumoto H, Hashimoto O, Nanba E, Nishida H, Sugiyama T, Kasai K, Watanabe K, Kano Y, Kato N, Sasaki T. Tachykinin 1 (TAC1) gene SNPs and haplotypes with autism: a case-control study. *Brain Dev.* 2007 Sep;29(8):510-3

Corcoran KE, Trzaska KA, Fernandes H, Bryan M, Taborga M, Srinivas V, Packman K, Patel PS, Rameshwar P. Mesenchymal stem cells in early entry of breast cancer into bone marrow. *PLoS One.* 2008 Jun 25;3(6):e2563

Howard MR, Millward-Sadler SJ, Vasilliou AS, Salter DM, Quinn JP. Mechanical stimulation induces preprotachykinin gene expression in osteoarthritic chondrocytes which is correlated with modulation of the transcription factor neuron restrictive silence factor. *Neuropeptides.* 2008 Oct-Dec;42(5-6):681-6

Murthy RG, Greco SJ, Taborga M, Patel N, Rameshwar P. Tac1 regulation by RNA-binding protein and miRNA in bone marrow stroma: Implication for hematopoietic activity. *Brain Behav Immun.* 2008 May;22(4):442-50

Rosso M, Robles-Frías MJ, Coveñas R, Salinas-Martín MV, Muñoz M. The NK-1 receptor is expressed in human primary gastric and colon adenocarcinomas and is involved in the antitumor action of L-733,060 and the mitogenic action of substance P on human gastrointestinal cancer cell lines. *Tumour Biol.* 2008;29(4):245-54

Chernova I, Lai JP, Li H, Schwartz L, Tuluc F, Korchak HM, Douglas SD, Kilpatrick LE. Substance P (SP) enhances CCL5-induced chemotaxis and intracellular signaling in human monocytes, which express the truncated neurokinin-1 receptor (NK1R). *J Leukoc Biol.* 2009 Jan;85(1):154-64

David S, Kan T, Cheng Y, Agarwal R, Jin Z, Mori Y. Aberrant silencing of the endocrine peptide gene tachykinin-1 in gastric cancer. *Biochem Biophys Res Commun.* 2009 Jan 16;378(3):605-9

Ellsworth RE, Seebach J, Field LA, Heckman C, Kane J, Hooke JA, Love B, Shriver CD. A gene expression signature that defines breast cancer metastases. *Clin Exp Metastasis.* 2009;26(3):205-13

Reddy BY, Greco SJ, Patel PS, Trzaska KA, Rameshwar P. RE-1-silencing transcription factor shows tumor-suppressor functions and negatively regulates the oncogenic TAC1 in breast cancer cells. *Proc Natl Acad Sci U S A.* 2009 Mar 17;106(11):4408-13

This article should be referenced as such:

Lim PK, Patel SA, Rameshwar P. TAC1 (tachykinin, precursor 1). *Atlas Genet Cytogenet Oncol Haematol.* 2010; 14(5):485-487.
