

# Gene Section

## Review

# GPX1 (Glutathione Peroxidase 1)

Mikhail V Kulak, Ronald J Weigel

Department of Surgery, University of Iowa, 200 Hawkins Drive, 1509 JCP, Iowa City, IA, 52242, United States (MVK, RJW)

Published in Atlas Database: November 2013

Online updated version : <http://AtlasGeneticsOncology.org/Genes/GPX1ID40747ch3p21.html>  
DOI: 10.4267/2042/53966

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.  
© 2014 Atlas of Genetics and Cytogenetics in Oncology and Haematology

## Abstract

Review on GPX1, with data on DNA/RNA, on the protein encoded and where the gene is implicated.

## Identity

**Other names:** GPXD, GSHPX1

**HGNC (Hugo):** GPX1

**Location:** 3p21.31

### Local order

- C3orf62, chr3:49306030-49314508, chromosome 3 open reading frame 62
- USP4, 3p21.31, chr3:49314577-49377536, ubiquitin specific peptidase 4 (proto-oncogene), transcript variant 2
- USP4, 3p21.31, chr3:49314577-49377536, ubiquitin specific peptidase 4 (proto-oncogene), transcript variant 1
- USP4, 3p21.31, chr3:49349218-49377536, ubiquitin specific peptidase 4 (proto-oncogene), transcript variant 3

- **GPX1**, 3p21.31, chr3:49394609-49395791, glutathione peroxidase 1, transcript variant 1
- **GPX1**, 3p21.31, chr3:49394609-49395791, glutathione peroxidase 1, transcript variant 2
- RHOA, 3p21.31, chr3:49396579-49449526, ras homolog family member A
- TCTA, 3p21.31, chr3:49449639-49453909, T-cell leukemia translocation altered
- AMT, 3p21.31, chr3:49454211-49460111, aminomethyltransferase

The colocalization of genes is presented on figure 1.


### Note

According to hg19/GRCh37-Feb\_2009:

- GPX1 at chr3:49394609-49395791 - (NM\_000581) glutathione peroxidase 1 isoform 1
- GPX1 at chr3:49394609-49395791 - (NM\_201397) glutathione peroxidase 1 isoform 2

GPX1 is an enzyme that reduces hydrogen and lipid peroxides to water or alcohols by reducing glutathione. Systematic name: glutathione: hydrogen-peroxide oxidoreductase.

Reaction: 2 glutathione + H<sub>2</sub>O<sub>2</sub> = glutathione disulfide + 2 H<sub>2</sub>O [RN:R00274].






Figure 2

## DNA/RNA

### Description

According to hg19/GRCh37-Feb\_2009:  
 - Start: chr3:49394609 bp from pter  
 - End: chr3:49395791 bp from pter  
 - Size: 1183 bases  
 - Orientation: minus strand

### Transcription

Two alternatively spliced transcript variants encoding distinct isoforms have been shown for this gene (Nucleotide).

Isoform 1 represents the shorter transcript (921 bases), RefSeq: NM\_000581.2, which is comprised of 2 exons and coding the longer isoform (Figure 2A).

Isoform 2 is 1200 bases, RefSeq: NM\_201397.1, also this variant is intronless.

Due to the fact that this variant is not spliced the open reading frame is shifted and the protein is shorter from C-terminus compared to isoform 1 (Figure 2B).

### Pseudogene

Two pseudogenes have been found so far. GPX1P1 glutathione peroxidase pseudogene 1 (other names: GPXL2, GPXP1) is located at the locus Xp22.2 (HGNC:4560). GPX1P2 glutathione peroxidase pseudogene 2 (other names: GPXP2, GPXP2P) is located at the locus 21q21.3 (HGNC:4561).

## Protein

### Description

203 aa (Accession: NM\_000581.2) isoform 1; 98 aa (Accession: NM\_201397.1) isoform 2.

GPX1 belongs to the family of glutathione peroxidases (Kryukov et al., 2003).

GPX1, GPX2, GPX3, GPX4 and GPX6 utilize a UGA codon that specifies insertion of a selenocysteine residue which is critical to protein function (Arthur, 2000). Both isoforms contain selenocysteine at the position 49 (Mullenbach et al., 1988).

### Expression

GPX1 is found at high levels in tissues exposed to high oxygen tensions such as in the lungs, at the cellular elements of blood, liver, kidney and pancreas, and also at moderate levels in heart, muscle and brain (Esposito et al., 2000; Moscow et al., 1992).

**Regulation:** Aberrant promoter methylation and consequence silencing has been shown for GPX1 expression during several pathological conditions in breast (Kulak et al., 2012) and gastric cancer (Min et al., 2012); whereas, induction of GPX1 gene expression was associated with transcription factors such TFAP2C in breast cancer (Kulak et al., 2012) and ZNF143 transcription factor under the mitochondrial respiratory dysfunction (Lu et al., 2012).

## Localisation

The protein is detected in cytoplasm and mitochondria but the ratio may vary and be dependent on cellular function (Chiu et al., 1976; Timcenko-Youssef et al., 1985; Reeves et al., 2009).

## Function

GPX1 is an enzyme of mammals and birds which protects against the damaging effects of various endogenously formed hydroperoxides and hydrogen peroxide as follows:  $H_2O_2 + 2\text{GSH} - 2\text{H}_2\text{O} + \text{GSS}$  and  $\text{ROOH} + 2\text{GSH} - \text{GSSG} + \text{ROH} + \text{H}_2\text{O}$  where ROOH represents lipid hydroperoxides, membrane associated phospholipid hydroperoxides (Ursini et al., 1985; Reeves et al., 2009).

## Homology

The GPX1 gene is present in vertebrates and across all mammals demonstrates homology of about 70% at the nucleotide level (Mariotti et al., 2012).

## Mutations

### Note

Single nucleotide and ALA polymorphism have been shown for GPX1.

The 5'-UTR of GPX1 was found to contain a single nucleotide C/T polymorphism rs 1800668 located at the position ch3:49395757. The CC genotype demonstrates relatively high activity of GPX1 compared to CT or TT variant of alleles; however, the polymorphism does not alter the protein structure so differences in activity might be associated with transcriptional regulation since rs 1080668 site is located within promoter of GPX1 (Najafi et al., 2012).

A single nucleotide polymorphism rs 1050450 at the position ch3:49394834 leads to Pro198->Leu (C->T) substitution in GPX1. Many publications suggested an associated increased risk of cancer with the TT (Leu) genotype.

However, extended meta-analysis failed to find a significant correlation of the polymorphism (rs 1050450) with cancer risk, although the TT GPX1 genotype examined in erythrocytes demonstrated significantly lower functional activity (Hong et al., 2012).

ALA N-terminal polymorphism has been shown for the GPX1 gene. In this variant, the number of GCG repeats is altered resulting in a protein with a variable number of alanines 5, 6, or 7 (Shen et al., 1994).

Some association between 5-ALA genotype and high risk of breast cancer have been shown (Knight et al., 2004). However, associations between specific genotypes and the risk of prostate cancer have not been found (Kote-Jarai et al., 2002).

## Implicated in

### Various cancers

#### Oncogenesis

GPX1 involvement in certain cancers has been shown by several line of evidence.

#### Lung cancer

##### Note

The expression of GPX1 may be altered through LOH (loss of heterozygosity) of GPX1 which leads to decrease gene activity and increased risk of lung cancer (Hardie et al., 2000). Up-regulation of GPX1 in erythrocytes may be seen under several conditions such as hypoxia or with treatment with chemicals such as alcohol (Raaschou-Nielsen et al., 2006) but GPX1 activity tended to be significantly lower in smokers compared to non-smokers (Ravn-Haren et al., 2006) and these differences have been hypothesized to account for increased risk of lung cancer in smokers (Ratnasinghe et al., 2000). However high level of GPX1 was shown in lung cancer compared to non-malignant tissue (Blomquist et al., 2009).

#### Breast cancer

##### Note

Decrease GPX1 activity due to LOH (Hu et al., 2003; Hu et al., 2005) or aberrant hypermethylation of the GPX1 gene promotor (Kulak et al., 2012) leads to high risk of breast cancer.

#### Renal cancer

##### Note

Selenium consumption plays a dramatic role on GPX1 activity. Selenium in the diet altered both the mRNA and protein levels of GPX1 in mice (Sunde et al., 2009) whereas in pigs selenoprotein gene expression and/or protein production is not dependent on prolonged dietary selenium deficiency or excess (Liu et al., 2012). In humans no association was found between the Se status and breast or colorectal cancer risk (Dennert et al., 2011), but poor Se status was indicative of high mortality rate in renal cancer patients (Jean-Claude et al., 2012; Meyer et al., 2012).

#### Gastric cancer

##### Note

Aberrant hypermethylation of the GPX1 gene promotor may decrease GPX1 expression and has been described as a mechanism of GPX1 silencing in gastric cancer (Jee et al., 2009).

#### Bladder cancer

##### Note

Relatively high level of GPX1 was observed in bladder cancer (Reszka, 2012).

## Anticancer drug resistance

### Note

One hypothesis proposes that when damaged cells have progressed to a precancerous status, increased GPX1 activity may become procarcinogenic, presumably due to inhibition of hydroperoxide-mediated apoptosis (Chu et al., 2004) and may be responsible for antitumor drug resistance such as doxorubicin (Gouazé et al., 2001) which acts through increasing ROS products in cells (Wang et al., 2004).

## Genetic polymorphisms and cancer

### Note

As mentioned above, genetic polymorphisms have been reported for GPX1.

Investigations have attempted to demonstrate an association with cancer incidence, however, no consistent association has been found to date (Lei et al., 2007; Arsova-Sarafinovska et al., 2009; Erdem et al., 2012; Hong et al., 2012).

## References

- Chiu DT, Stults FH, Tappel AL. Purification and properties of rat lung soluble glutathione peroxidase. *Biochim Biophys Acta*. 1976 Oct 11;445(3):558-66
- Timcenko-Youssef L, Yamazaki RK, Kimura T. Subcellular localization of adrenal cortical glutathione peroxidase and protective role of the mitochondrial enzyme against lipid peroxidative damage. *J Biol Chem*. 1985 Oct 25;260(24):13355-9
- Ursini F, Maiorino M, Gregolin C. The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. *Biochim Biophys Acta*. 1985 Mar 29;839(1):62-70
- Mullenbach GT, Tabrizi A, Irvine BD, Bell GI, Tainer JA, Hallewell RA. Selenocysteine's mechanism of incorporation and evolution revealed in cDNAs of three glutathione peroxidases. *Protein Eng*. 1988 Sep;2(3):239-46
- Moscow JA, Morrow CS, He R, Mullenbach GT, Cowan KH. Structure and function of the 5'-flanking sequence of the human cytosolic selenium-dependent glutathione peroxidase gene (hgp1). *J Biol Chem*. 1992 Mar 25;267(9):5949-58
- Shen Q, Townes PL, Padden C, Newburger PE. An in-frame trinucleotide repeat in the coding region of the human cellular glutathione peroxidase (GPX1) gene: in vivo polymorphism and in vitro instability. *Genomics*. 1994 Sep 1;23(1):292-4
- Arthur JR. The glutathione peroxidases. *Cell Mol Life Sci*. 2000 Dec;57(13-14):1825-35
- Esposito LA, Kokoszka JE, Waymire KG, Cottrell B, MacGregor GR, Wallace DC. Mitochondrial oxidative stress in mice lacking the glutathione peroxidase-1 gene. *Free Radic Biol Med*. 2000 Mar 1;28(5):754-66
- Hardie LJ, Briggs JA, Davidson LA, Allan JM, King RF, Williams GI, Wild CP. The effect of hOGG1 and glutathione peroxidase I genotypes and 3p chromosomal loss on 8-hydroxydeoxyguanosine levels in lung cancer. *Carcinogenesis*. 2000 Feb;21(2):167-72
- Ratnasinghe D, Tangrea JA, Andersen MR, Barrett MJ, Virtamo J, Taylor PR, Albanes D. Glutathione peroxidase codon 198 polymorphism variant increases lung cancer risk. *Cancer Res*. 2000 Nov 15;60(22):6381-3
- Gouazé V, Mirault ME, Carpentier S, Salvayre R, Levade T, Andrieu-Abadie N. Glutathione peroxidase-1 overexpression prevents ceramide production and partially inhibits apoptosis in doxorubicin-treated human breast carcinoma cells. *Mol Pharmacol*. 2001 Sep;60(3):488-96
- Kote-Jarai Z, Durocher F, Edwards SM, Hamoudi R, Jackson RA, Ardern-Jones A, Murkin A, Dearnaley DP, Kirby R, Houlston R, Easton DF, Eeles R. Association between the GCG polymorphism of the selenium dependent GPX1 gene and the risk of young onset prostate cancer. *Prostate Cancer Prostatic Dis*. 2002;5(3):189-92
- Hu YJ, Diamond AM. Role of glutathione peroxidase 1 in breast cancer: loss of heterozygosity and allelic differences in the response to selenium. *Cancer Res*. 2003 Jun 15;63(12):3347-51
- Kryukov GV, Castellano S, Novoselov SV, Lobanov AV, Zehab O, Guigó R, Gladyshev VN. Characterization of mammalian selenoproteomes. *Science*. 2003 May 30;300(5624):1439-43
- Chu FF, Esworthy RS, Doroshow JH. Role of Selenium-dependent glutathione peroxidases in gastrointestinal inflammation and cancer. *Free Radic Biol Med*. 2004 Jun 15;36(12):1481-95
- Knight JA, Onay UV, Wells S, Li H, Shi EJ, Andrusis IL, Ozcelik H. Genetic variants of GPX1 and SOD2 and breast cancer risk at the Ontario site of the Breast Cancer Family Registry. *Cancer Epidemiol Biomarkers Prev*. 2004 Jan;13(1):146-9
- Wang S, Konorev EA, Kotamraju S, Joseph J, Kalivendi S, Kalyanaraman B. Doxorubicin induces apoptosis in normal and tumor cells via distinctly different mechanisms. Intermediacy of H<sub>2</sub>O<sub>2</sub>- and p53-dependent pathways. *J Biol Chem*. 2004 Jun 11;279(24):25535-43
- Hu Y, Benya RV, Carroll RE, Diamond AM. Allelic loss of the gene for the GPX1 selenium-containing protein is a common event in cancer. *J Nutr*. 2005 Dec;135(12 Suppl):3021S-3024S
- Ravn-Haren G, Olsen A, Tjønneland A, Dragsted LO, Nexø BA, Wallin H, Overvad K, Raaschou-Nielsen O, Vogel U. Associations between GPX1 Pro198Leu polymorphism, erythrocyte GPX activity, alcohol consumption and breast cancer risk in a prospective cohort study. *Carcinogenesis*. 2006 Apr;27(4):820-5
- Lei XG, Cheng WH, McClung JP. Metabolic regulation and function of glutathione peroxidase-1. *Annu Rev Nutr*. 2007;27:41-61
- Arsova-Sarafinovska Z, Matevska N, Eken A, Petrovski D, Banev S, Dzikova S, Georgiev V, Sikole A, Erdem O, Sayal A, Aydin A, Dimovski AJ. Glutathione peroxidase 1 (GPX1) genetic polymorphism, erythrocyte GPX activity, and prostate cancer risk. *Int Urol Nephrol*. 2009;41(1):63-70
- Blomquist T, Crawford EL, Mullins D, Yoon Y, Hernandez DA, Khuder S, Ruppel PL, Peters E, Oldfield DJ, Austermiller B, Anders JC, Willey JC. Pattern of antioxidant and DNA repair gene expression in normal airway epithelium associated with lung cancer diagnosis. *Cancer Res*. 2009 Nov 15;69(22):8629-35
- Jee CD, Kim MA, Jung EJ, Kim J, Kim WH. Identification of

- genes epigenetically silenced by CpG methylation in human gastric carcinoma. *Eur J Cancer*. 2009 May;45(7):1282-93
- Reeves MA, Hoffmann PR. The human selenoproteome: recent insights into functions and regulation. *Cell Mol Life Sci*. 2009 Aug;66(15):2457-78
- Sunde RA, Raines AM, Barnes KM, Evenson JK. Selenium status highly regulates selenoprotein mRNA levels for only a subset of the selenoproteins in the selenoproteome. *Biosci Rep*. 2009 Jun 25;29(5):329-38
- Dennert G, Zwahlen M, Brinkman M, Vinceti M, Zeegers MP, Horneber M. Selenium for preventing cancer. *Cochrane Database Syst Rev*. 2011 May 11;(5):CD005195
- Erdem O, Eken A, Akay C et al.. Association of GPX1 polymorphism, GPX activity and prostate cancer risk. *Hum Exp Toxicol*. 2012 Jan;31(1):24-31
- Jean-Claude M, Emmanuelle P, Juliette H, Michèle B, Gérard D, Eric F, Xavier H, Bertrand L, Jean-Fabien Z, Yves P, Gérard N. Clinical and economic impact of malnutrition per se on the postoperative course of colorectal cancer patients. *Clin Nutr*. 2012 Dec;31(6):896-902
- Liu Y, Zhao H, Zhang Q, Tang J, Li K, Xia XJ, Wang KN, Li K, Lei XG. Prolonged dietary selenium deficiency or excess does not globally affect selenoprotein gene expression and/or protein production in various tissues of pigs. *J Nutr*. 2012 Aug;142(8):1410-6
- Lu W, Chen Z, Zhang H, Wang Y, Luo Y, Huang P. ZNF143 transcription factor mediates cell survival through upregulation of the GPX1 activity in the mitochondrial respiratory dysfunction. *Cell Death Dis*. 2012 Nov 15;3:e422
- Mariotti M, Ridge PG, Zhang Y, Lobanov AV, Pringle TH, Guigo R, Hatfield DL, Gladyshev VN. Composition and evolution of the vertebrate and mammalian selenoproteomes. *PLoS One*. 2012;7(3):e33066
- Meyer HA, Endermann T et al.. Selenoprotein P status correlates to cancer-specific mortality in renal cancer patients. *PLoS One*. 2012;7(10):e46644
- Min SY, Kim HS, Jung EJ, Jung EJ, Jee CD, Kim WH. Prognostic significance of glutathione peroxidase 1 (GPX1) down-regulation and correlation with aberrant promoter methylation in human gastric cancer. *Anticancer Res*. 2012 Aug;32(8):3169-75
- Najafi M, Ghasemi H, Roustazadeh A, Alipoor B. Phenotype and genotype relationship of glutathione peroxidase1 (GPx1) and rs 1800668 variant: the homozygote effect on kinetic parameters. *Gene*. 2012 Aug 15;505(1):19-22
- Reszka E. Selenoproteins in bladder cancer. *Clin Chim Acta*. 2012 May 18;413(9-10):847-54
- Hong Z, Tian C, Zhang X. GPX1 gene Pro200Leu polymorphism, erythrocyte GPX activity, and cancer risk. *Mol Biol Rep*. 2013 Feb;40(2):1801-12
- Kulak MV, Cyr AR, Woodfield GW, Bogachek M, Spanheimer PM, Li T, Price DH, Domann FE, Weigel RJ. Transcriptional regulation of the GPX1 gene by TFAP2C and aberrant CpG methylation in human breast cancer. *Oncogene*. 2013 Aug 22;32(34):4043-51
- 
- This article should be referenced as such:*
- Kulak MV, Weigel RJ. GPX1 (Glutathione Peroxidase 1). *Atlas Genet Cytogenet Oncol Haematol*. 2014; 18(7):465-469.
-