

Gene Section Review

FOXQ1 (forkhead box Q1)

Jon Christensen, Pascale Anderle

Institute for Macromolecular Chemistry, Alber-Ludwigs-University of Freiburg, Freiburg, Germany (JC), Swiss Institute of Bioinformatics, Lausanne, Switzerland (PA)

Published in Atlas Database: April 2014

Online updated version : <http://AtlasGeneticsOncology.org/Genes/FOXQ1ID45906ch6p25.html>
DOI: 10.4267/2042/55371

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 2015 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

Review on FOXQ1, with data on DNA/RNA, on the protein encoded and where the gene is implicated.

Identity

Other names: HFH1

HGNC (Hugo): FOXQ1

Location: 6p25.3

DNA/RNA

Description

The FOXQ1 gene is 2338 base pairs in length and is intronless.

Transcription

In mouse Foxq1 has been reported to be regulated by Hoxa1 (Martinez-Ceballos et al., 2005), Hoxc13 (Potter et al., 2006) and Tgf β (Zhang et al., 2011). In human FOXQ1 has been shown to be a target of the Wnt pathway (Christensen et al., 2013; Xia et al., 2014).

Protein

Description

The forkhead box Q1 gene codes for a 403 amino acid long protein with a size of 41.5 kDa. FOXQ1

is a member of the Forkhead box (Fox) superfamily.




Figure 1. Structure of the winged helix domain. α -helices are shown as red cylinders (H1, H2 and H3), β -strands as blue arrows and W1 and W2 denote the wings (Clark et al., 1993; Gajiwala and Burley, 2000).

The family members share a conserved DNA-binding domain named forkhead box or winged helix domain.

The domain consists of three α -helices, three β -sheets and two loops termed wings.

Expression

Predominantly in the stomach, trachea, bladder and salivary gland (Bieller et al., 2001).




Figure 2. Graphical illustration of the FOXQ1 amino acid sequence and domains. (I) Acid and serine-rich domain. (WH) winged helix domain. (II) Serin-rich domain. (III) Proline-rich domain. (IV) Functional conserved domain (Hong et al., 2001; Wu et al., 2013).

Function

In mice *Foxq1* is involved in hair follicle differentiation (Hong et al., 2001; Potter et al., 2006).

A mutation in the *Foxq1* gene is responsible for an impaired differentiation of the hair shaft in the satin mice (Hong et al., 2001). In the digestive system *Foxq1* has been shown to regulate acid secretion and expression of *Muc5ac* (Goering et al., 2008; Verzi et al., 2008).

Homology

According to NCBI the following genes have been suggested to be putative homologues: FOXQ1 (*H. sapiens*), *Foxq1* (*M. musculus*), *Foxq1* (*R. norvegicus*), *Foxq1a* (*D. rerio*) and *Foxq1b* (*D. rerio*). Conserved domains from CDD found in protein sequences by rpsblast searching was FH (c100061).

Mutations

Note

Mutations in the *Foxq1* gene is responsible for the hair follicle defects seen in the satin mouse mutant. Three mutations have been described leading to similar phenotypes of the animals. *Foxq1^{sa}* has a 67 bp deletion from 686-752 and a base pair change CA-AT at position 766-767. *Foxq1^{sa-el}* has a point mutation a position 383 changing T to G thus replacing isoleucine with serin at position 128 in the protein. *Foxq1^{sa-j}* has C to T mutation in position 490 changing the amino acid arginine to cysteine at position 164 in the protein (Hong et al., 2001; Wu et al., 2013).

Implicated in

Bladder cancer

Oncogenesis

FOXQ1 was overexpressed in bladder cancer samples. Depletion of FOXQ1 expression in bladder cancer cell lines reduced invasiveness and EMT markers (Zhu et al., 2013).

Breast cancer

Prognosis

FOXQ1 expression in breast cancer patients is associated with poor survival, high grade, metastatic status and basal-like phenotype (Qiao et al., 2011).

Oncogenesis

FOXQ1 overexpression was observed in invasive breast cancer cell lines compared to non-invasive. FOXQ1 expression increases breast cancer cell proliferation, migration and invasion in vitro and metastasis in vivo (Zhang et al., 2011). FOXQ1 promotes an EMT phenotype through transcriptional regulation of *CDH1* (Qiao et al., 2011; Zhang et al., 2011).

Colorectal cancer

Oncogenesis

Several studies have shown FOXQ1 to be overexpressed in colorectal tumor samples compared to healthy colonocytes (Bieller et al., 2001; Sabates-Bellver et al., 2007; Kaneda et al., 2010; Christensen et al., 2013).

The increased expression of FOXQ1 could be due to a hyperactive Wnt pathway in these tumors. Wnt activity directly correlates with FOXQ1 expression in colorectal cancer cell lines and β -catenin can bind to the promoter of FOXQ1 and increase transcription (Christensen et al., 2013). FOXQ1 expression can induce an EMT phenotype (Qiao et al., 2011; Abba et al., 2013). FOXQ1 does not increase growth but seems to protect from apoptosis (Kaneda et al., 2010; Qiao et al., 2011; Abba et al., 2013). The anti-apoptotic effect was mediated through FOXQ1 regulation of p21 (Kaneda et al., 2010).

Gastric cancer

Prognosis

The expression of FOXQ1 was a prognostic factor for overall survival and correlated with tumor size, grade and tumor-node metastasis stage (Liang et al., 2013).

Oncogenesis

FOXQ1 increases migration and proliferation by downregulating NRXN3.

Glioma**Disease**

Tumors that arise from the glial cells, the most common site is the brain.

Oncogenesis

FOXQ1 increased migration and proliferation by downregulating NRXN3 (Sun et al., 2013).

Hepatocarcinoma**Cytogenetics**

FOXQ1 correlated with overall worse survival and higher recurrence (Wang et al., 2013; Xia et al., 2014).

Oncogenesis

In hepatocarcinoma FOXQ1 directly activated the EMT transcription factor ZEB2. This led to an EMT phenotype and increased lung metastasis. FOXQ1 and ZEB2 expression correlated positively in hepatocarcinoma samples but inversely with CDH1. FOXQ1 induced metastasis through regulation of VersicanV1, which promoted tumor-associated-macrophages attraction. Also, similarly to colorectal cancer expression of FOXQ1 was regulated by the Wnt pathway in hepatocarcinoma (Xia et al., 2014).

Non-small-cell lung carcinoma**Prognosis**

FOXQ1 expression was associated with poor prognosis and EMT (Feng et al., 2012).

Ovarian cancer**Oncogenesis**

FOXQ1 expression increased ovarian cancer cell proliferation, invasion and induced an EMT phenotype (Gao et al., 2012).

References

Clark KL, Halay ED, Lai E, Burley SK. Co-crystal structure of the HNF-3/fork head DNA-recognition motif resembles histone H5. *Nature*. 1993 Jul 29;364(6436):412-20

Gajiwala KS, Burley SK. Winged helix proteins. *Curr Opin Struct Biol*. 2000 Feb;10(1):110-6

Bieller A, Pasche B, Frank S, Gläser B, Kunz J, Witt K, Zoll B. Isolation and characterization of the human forkhead gene FOXQ1. *DNA Cell Biol*. 2001 Sep;20(9):555-61

Hong HK, Noveroske JK, Headon DJ, Liu T, Sy MS, Justice MJ, Chakravarti A. The winged helix/forkhead transcription factor Foxq1 regulates differentiation of hair in satin mice. *Genesis*. 2001 Apr;29(4):163-71

Martinez-Ceballos E, Chambon P, Gudas LJ. Differences in gene expression between wild type and Hoxa1 knockout embryonic stem cells after retinoic acid treatment or leukemia inhibitory factor (LIF) removal. *J Biol Chem*. 2005 Apr 22;280(16):16484-98

Potter CS, Peterson RL, Barth JL, Pruett ND, Jacobs DF, Kern MJ, Argraves WS, Sundberg JP, Awgulewitsch A. Evidence that the satin hair mutant gene Foxq1 is among multiple and functionally diverse regulatory targets for Hoxc13 during hair follicle differentiation. *J Biol Chem*. 2006 Sep 29;281(39):29245-55

Sabates-Bellver J, Van der Flier LG, de Palo M, Cattaneo E, Maake C, Rehrauer H, Laczko E, Kurowski MA, Bujnicki JM, Menigatti M, Luz J, Ranalli TV, Gomes V, Pastorelli A, Faggiani R, Anti M, Jiricny J, Clevers H, Marra G. Transcriptome profile of human colorectal adenomas. *Mol Cancer Res*. 2007 Dec;5(12):1263-75

Goering W, Adham IM, Pasche B, Manner J, Ochs M, Engel W, Zoll B. Impairment of gastric acid secretion and increase of embryonic lethality in Foxq1-deficient mice. *Cytogenet Genome Res*. 2008;121(2):88-95

Verzi MP, Khan AH, Ito S, Shivdasani RA. Transcription factor foxq1 controls mucin gene expression and granule content in mouse stomach surface mucous cells. *Gastroenterology*. 2008 Aug;135(2):591-600

Kaneda H, Arao T, Tanaka K, Tamura D, Aomatsu K, Kudo K, Sakai K, De Velasco MA, Matsumoto K, Fujita Y, Yamada Y, Tsurutani J, Okamoto I, Nakagawa K, Nishio K. FOXQ1 is overexpressed in colorectal cancer and enhances tumorigenicity and tumor growth. *Cancer Res*. 2010 Mar 1;70(5):2053-63

Feuerborn A, Srivastava PK, Küffer S, Grandy WA, Sijmonsma TP, Gretz N, Brors B, Gröne HJ. The Forkhead factor FoxQ1 influences epithelial differentiation. *J Cell Physiol*. 2011 Mar;226(3):710-9

Qiao Y, Jiang X, Lee ST, Karuturi RK, Hooi SC, Yu Q. FOXQ1 regulates epithelial-mesenchymal transition in human cancers. *Cancer Res*. 2011 Apr 15;71(8):3076-86

Zhang H, Meng F, Liu G, Zhang B, Zhu J, Wu F, Ethier SP, Miller F, Wu G. Forkhead transcription factor foxq1 promotes epithelial-mesenchymal transition and breast cancer metastasis. *Cancer Res*. 2011 Feb 15;71(4):1292-301

Feng J, Zhang X, Zhu H, Wang X, Ni S, Huang J. FoxQ1 overexpression influences poor prognosis in non-small cell lung cancer, associates with the phenomenon of EMT. *PLoS One*. 2012;7(6):e39937

Gao M, Shih leM, Wang TL. The role of forkhead box q1 transcription factor in ovarian epithelial carcinomas. *Int J Mol Sci*. 2012 Oct 26;13(11):13881-93

Abba M, Patil N, Rasheed K, Nelson LD, Mudduluru G, Leupold JH, Allgayer H. Unraveling the role of FOXQ1 in colorectal cancer metastasis. *Mol Cancer Res*. 2013 Sep;11(9):1017-28

Christensen J, Bentz S, Sengstag T, Shastri VP, Anderle P. FOXQ1, a novel target of the Wnt pathway and a new marker for activation of Wnt signaling in solid tumors. *PLoS One*. 2013;8(3):e60051

Liang SH, Yan XZ, Wang BL, Jin HF, Yao LP, Li YN, Chen M, Nie YZ, Wang X, Guo XG, Wu KC, Ding J, Fan DM. Increased expression of FOXQ1 is a prognostic marker for patients with gastric cancer. *Tumour Biol*. 2013 Oct;34(5):2605-9

Sun HT, Cheng SX, Tu Y, Li XH, Zhang S. FoxQ1 promotes glioma cells proliferation and migration by regulating NRXN3 expression. *PLoS One*. 2013;8(1):e55693

Wang W, He S, Ji J, Huang J, Zhang S, Zhang Y. The prognostic significance of FOXQ1 oncogene

overexpression in human hepatocellular carcinoma. *Pathol Res Pract*. 2013 Jun;209(6):353-8

Wu B, Herbert Pratt C, Potter CS, Silva KA, Kennedy V, Sundberg JP. R164C mutation in FOXQ1 H3 domain affects formation of the hair medulla. *Exp Dermatol*. 2013 Mar;22(3):234-6

Zhu Z, Zhu Z, Pang Z, Xing Y, Wan F, Lan D, Wang H. Short hairpin RNA targeting FOXQ1 inhibits invasion and metastasis via the reversal of epithelial-mesenchymal transition in bladder cancer. *Int J Oncol*. 2013

Apr;42(4):1271-8

Xia L, Huang W, Tian D, Zhang L, Qi X, Chen Z, Shang X, Nie Y, Wu K. Forkhead box Q1 promotes hepatocellular carcinoma metastasis by transactivating ZEB2 and VersicanV1 expression. *Hepatology*. 2014 Mar;59(3):958-73

This article should be referenced as such:

Christensen J, Anderle P. FOXQ1 (forkhead box Q1). *Atlas Genet Cytogenet Oncol Haematol*. 2015; 19(1):11-14.
