

Gene Section

Review

PPP2R1A (protein phosphatase 2 regulatory subunit A, alpha)

Razia Sultana, Kentaro Nakayama, Kohei Nakamura, Satoru Kyo

Department of Gynecology/Obstetrics Shimane University Hospital kn88@med.shimane-u.ac.jp

Published in Atlas Database: March 2016

Online updated version : <http://AtlasGeneticsOncology.org/Genes/PPP2R1AID41814ch19q13.html>

Printable original version : <http://documents.irevues.inist.fr/bitstream/handle/2042/66936/03-2016-PPP2R1AID41814ch19q13.pdf>

DOI: 10.4267/2042/66936

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2016 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

Review on PPP2R1A, with data on DNA, on the protein encoded, and where the gene is implicated.

Identity

Other names: 2AAA, MGC786, PP2A-Aalpha, PP2AAALPHA, PR65A

HGNC (Hugo): PPP2R1A

Location: 19q13.41

Local order: Start at 52693055 and end at 52729678bp. (NCBI 37, August 2010), on the direct strand.

DNA/RNA

Description

PPP2R1A gene is encoded by 15 exons located on

chromosome 19q13. The genomic size is 36624 bp.

Transcription

mRNA size: 2509 bp; coding sequence from 296bp-1770 bp.

Protein


Note

The PPP2R1A cDNA has 1767 bp open reading frame encoding a predicted polypeptide of 589 amino acids with a predicted molecular mass of 65KDa.


Description


The PR65 subunit of protein phosphatase 2A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit.

Required for proper chromosome segregation and for centromeric localization of SGOL1 in mitosis.



PPP2R1A gene is located on the long (q) arm of chromosome 19 at position 13.41





Found in a complex with at least ARL2, PPP2CB, PPP2R1A, PPP2R2A, PPP2R5E and TBCD. PP2A consists of a common heterodimeric core enzyme, composed of PPP2CA a 36 kDa catalytic subunit (subunit C) and PPP2R1A a 65 kDa constant regulatory subunit (PR65 or subunit A), that associates with a variety of regulatory subunits. Proteins that associate with the core dimer include three families of regulatory subunits B (the R2/B/PR55/B55, R3/B"/PR72/PR130/PR59 and R5/B'/B56 families), the 48 kDa variable regulatory subunit, viral proteins, and cell signaling molecules. Interacts with IPO9. Interacts with TP53 and SGOL1. Interacts with PLA2G16; this interaction might decrease PP2A activity. Belongs to the phosphatase 2A regulatory subunit A family.

Expression

PPP2R1A is highly expressed from brain (seen 50 times), skin , eye , lung , ovary , uterus , leiomyosarcoma and 129 other tissues.

Localisation

PPP2R1A is located in the cytoplasm, chromosome and centromere in the cell.

Function

PP2A is an important and ubiquitously expressed serine threonine phosphatase family, which plays a critical role in many fundamental cellular processes, such as cell proliferation, signal transduction, DNA repair, and apoptosis. The basis for its multifunctionality rests on the large number of subunits that determine its phosphatase activity, substrate specificity, and subcellular localization. PP2A is a heterotrimer composed of a scaffold subunit (A), a catalytic subunit (C), and one of many regulatory subunits (B). The dimeric form consisting of the scaffold and catalytic subunit also exists as the core-PP2A enzyme. Among these subunits, PP2A-A acts as a structural assembly base to escort the catalytic subunit and to facilitate interaction with the regulatory subunit and other substrates, which is essential for the activity of the holoenzyme. The PP2A scaffold subunit is encoded by two distinct genes, Ppp2r1a and Ppp2r1b, resulting in two

isoforms, A α and A β , which are 87% identical. However, in about 90% of the PP2A assemblies, the core and/or holoenzyme is composed of the A α scaffold subunit and is highly abundant in all tissues. The discovery that A α is mutated in a variety of human malignancies, including cancer of the breast, lung, skin, and ovaries, provided evidence that PP2A-A α plays a role in cancer, suggesting its role in tumor suppression. PP2A has been studied in vitro using an exogenous inhibitor, okadaic acid(OA), or RNA interference by others, suggesting its possible role in the regulation of meiosis. Recently, Su et al. found that up-regulation of protein phosphatase 2 catalytic subunit (PPP2CB) is key to the meiotic arrest phenotype, which demonstrated that PP2A might be involved in the regulation of meiosis resumption. Qi et al. and Chambon et al. both found that I2PP2A, the inhibitor of PP2A, was essential for sister chromatid segregation, which also suggested the importance of PP2A in meiosis.


Homology

The PPP2R1A gene is conserved in chimpanzee, Rhesus monkey, dog, cow, mouse, rat, fruit fly, mosquito, C.elegans, S.cerevisiae, K.lactis, E.gossypii, S.pombe, M.oryzae, N.crassa, A.thaliana, and frog.

Mutations

Somatic

Recently, somatic mutations in PPP2R1A have been reported in certain types of ovarian and uterine carcinoma. PPP2R1A mutations were demonstrated in 10 of 110 type I ovarian tumors (9.1%) including low grade serous, low-grade endometrioid, clear-cell, and mucinous carcinomas. Moreover, PPP2R1A mutations were observed in 2 of 30 type I uterine (endometrioid) carcinomas (6.7%) and 5 of 26 type II uterine (serous) carcinomas (19.2%). All mutations were located in the alpha-helix repeats near the interface between the A subunit and the regulatory B subunit of the enzyme complex. All mutations identified were heterozygous missense mutations.



Structure of PP2A: C is the catalytic subunit, A is the second regulatory or structural subunit, and B/B'/B''/B''' are the third variable subunits, which are structurally unrelated. In Mammalia, A and C are encoded by two genes (α and β); the B/PR55 subunits are encoded by four related genes (α , β , γ , and δ); the B'/PR61 family are encoded by five related genes (α , β , γ , δ and ϵ), some of which give rise to alternatively spliced products; the B'' family probably contains three related genes, encoding PR48, PR59 and the splice variants PR72 and PR130; SG2NA and striatin comprise the B''' subunit family

Implicated in

Wilms tumorigenesis

Note

PP2A plays a role in the Wnt signaling pathway and has been shown to be present in the multiprotein complex containing the adenomatous polyposis coli (APC) protein, the serine/threonine kinase GSK3B, and AXIN1 (axin) (Fagotto et al., 1999; Hsu et al., 1999; Seeling et al., 1999; Ikeda et al., 2000), suggesting that PP2A has a role in regulating the activity of this complex. Overexpression of the PP2A B56 regulatory subunit has been shown to markedly reduce CTNNB1 levels in mammalian cells (Seeling et al., 1999), which suggests that increased levels of PP2A accelerate the phosphorylation-dependent proteolysis of CTNNB1. Interestingly, CTNNB1 mutations have recently been reported to occur in about 15% of Wilms tumors (Koesters et al., 1999; Maiti et al., 2000), suggesting that the APC/CTNNB1/HNF4A (Tcf)-LEF1 pathway is involved in Wilms tumorigenesis. Since PP2A also plays a role in the Wnt signaling pathway, alterations in one or more of the PP2A subunits could plausibly affect this pathway and contribute to Wilms tumorigenesis.

Mental retardation, autosomal dominant 36 (MRD36)

Note

A form of mental retardation, a disorder characterized by significantly below average general intellectual functioning associate with impairments in adaptive behavior and manifested during the developmental period. The disease is identified 2 different de novo heterozygous missense mutation in

the PPP2R1A gene (Hougo et al., 2015). The mutation were found by parent-child trio exome sequencing and confirmed by Sanger sequencing. In vitro functional expression study showed that the patients with MRD36, PPP2R1A mutation affected PP2A holoenzyme formation by variably interfering with interaction of the A-alpha subunit with the C subunit. All mutation resulted in decreased phosphatase activity, consisting with a dominant negative effect.

References

- Alexandre H, Van Cauwenberge A, Tsukitani Y, Mulnard J. Pleiotropic effect of okadaic acid on maturing mouse oocytes. *Development*. 1991 Aug;112(4):971-80
- Calin GA, di lasio MG, Caprini E, Vorechovsky I, Natali PG, Sozzi G, Croce CM, Barbanti-Brodano G, Russo G, Negrini M. Low frequency of alterations of the alpha (PPP2R1A) and beta (PPP2R1B) isoforms of the subunit A of the serine-threonine phosphatase 2A in human neoplasms. *Oncogene*. 2000 Feb 24;19(9):1191-5
- Chambon JP, Touati SA, Berneau S, Cladière D, Hebras C, Groeme R, McDougall A, Wassmann K. The PP2A inhibitor I2PP2A is essential for sister chromatid segregation in oocyte meiosis II. *Curr Biol*. 2013 Mar 18;23(6):485-90
- Fagotto F, Jho Eh, Zeng L, Kurth T, Joos T, Kaufmann C, Costantini F. Domains of axin involved in protein-protein interactions, Wnt pathway inhibition, and intracellular localization *J Cell Biol* 1999 May 17;145(4):741-56
- Hemmings BA, Adams-Pearson C, Maurer F, Müller P, Goris J, Merlevede W, Hofsteenge J, Stone SR. alpha- and beta-forms of the 65-kDa subunit of protein phosphatase 2A have a similar 39 amino acid repeating structure *Biochemistry* 1990 Apr 3;29(13):3166-73
- Hsu W, Zeng L, Costantini F. Identification of a domain of Axin that binds to the serine/threonine protein phosphatase 2A and a self-binding domain *J Biol Chem* 1999 Feb 5;274(6):3439-45

- Ikeda S, Kishida M, Matsuura Y, Usui H, Kikuchi A. GSK-3beta-dependent phosphorylation of adenomatous polyposis coli gene product can be modulated by beta-catenin and protein phosphatase 2A complexed with Axin *Oncogene* 2000 Jan 27;19(4):537-45
- Koesters R, Ridder R, Kopp-Schneider A, Betts D, Adams V, Niggli F, Briner J, von Knebel Doeberitz M. Mutational activation of the beta-catenin proto-oncogene is a common event in the development of Wilms' tumors *Cancer Res* 1999 Aug 15;59(16):3880-2
- Kremmer E, Ohst K, Kiefer J, Brewis N, Walter G. Separation of PP2A core enzyme and holoenzyme with monoclonal antibodies against the regulatory A subunit: abundant expression of both forms in cells *Mol Cell Biol* 1997 Mar;17(3):1692-701
- Lu Q, Dunn RL, Angeles R, Smith GD. Regulation of spindle formation by active mitogen-activated protein kinase and protein phosphatase 2A during mouse oocyte meiosis *Biol Reprod* 2002 Jan;66(1):29-37
- Mailhes JB, Hilliard C, Fuseler JW, London SN. Okadaic acid, an inhibitor of protein phosphatase 1 and 2A, induces premature separation of sister chromatids during meiosis I and aneuploidy in mouse oocytes in vitro *Chromosome Res* 2003;11(6):619-31
- Maiti S, Alam R, Amos CI, Huff V. Frequent association of beta-catenin and WT1 mutations in Wilms tumors *Cancer Res* 2000 Nov 15;60(22):6288-92
- Pandey P, Seshacharyulu P, Das S, Rachagani S, Ponnusamy MP, Yan Y, Johansson SL, Datta K, Fong Lin M, Batra SK. Impaired expression of protein phosphatase 2A subunits enhances metastatic potential of human prostate cancer cells through activation of AKT pathway *Br J Cancer* 2013 Jun 25;108(12):2590-600
- Price NE, Mumby MC. Effects of regulatory subunits on the kinetics of protein phosphatase 2A *Biochemistry* 2000 Sep 19;39(37):11312-8
- Qi ST, Wang ZB, Ouyang YC, Zhang QH, Hu MW, Huang X, Ge Z, Guo L, Wang YP, Hou Y, Schatten H, Sun QY. Overexpression of SET β , a protein localizing to centromeres, causes precocious separation of chromatids during the first meiosis of mouse oocytes *J Cell Sci* 2013 Apr 1;126(Pt 7):1595-603
- Ruediger R, Ruiz J, Walter G. Human cancer-associated mutations in the A α subunit of protein phosphatase 2A increase lung cancer incidence in A α knock-in and knockout mice *Mol Cell Biol* 2011 Sep;31(18):3832-44
- Sablina AA, Chen W, Arroyo JD, Corral L, Hector M, Bulmer SE, DeCaprio JA, Hahn WC. The tumor suppressor PP2A A β regulates the RalA GTPase *Cell* 2007 Jun 1;129(5):969-82
- Seeling JM, Miller JR, Gil R, Moon RT, White R, Virshup DM. Regulation of beta-catenin signaling by the B56 subunit of protein phosphatase 2A *Science* 1999 Mar 26;283(5410):2089-91
- Su YQ, Sugiura K, Sun F, Pendola JK, Cox GA, Handel MA, Schimenti JC, Eppig JJ. MARF1 regulates essential oogenic processes in mice *Science* 2012 Mar 23;335(6075):1496-9
- Sun QY, Wu GM, Lai L, Bonk A, Cabot R, Park KW, Day BN, Prather RS, Schatten H. Regulation of mitogen-activated protein kinase phosphorylation, microtubule organization, chromatin behavior, and cell cycle progression by protein phosphatases during pig oocyte maturation and fertilization in vitro *Biol Reprod* 2002 Mar;66(3):580-8
- Swain JE, Wang X, Saunders TL, Dunn R, Smith GD. Specific inhibition of mouse oocyte nuclear protein phosphatase-1 stimulates germinal vesicle breakdown *Mol Reprod Dev* 2003 May;65(1):96-103
- Zhou J, Pham HT, Ruediger R, Walter G. Characterization of the A α and A β subunit isoforms of protein phosphatase 2A: differences in expression, subunit interaction, and evolution *Biochem J* 2003 Jan 15;369(Pt 2):387-98
-
- This article should be referenced as such:*
- Sultana R, Nakayama K, Nakamura K, Kyo S. PPP2R1A (protein phosphatase 2 regulatory subunit A, alpha). *Atlas Genet Cytogenet Oncol Haematol*. 2016; 20(11):563-566.
-