

Gene Section

Short Communication

OCA2 (OCA2 melanosomal transmembrane protein)

Kunal Ray, Mainak Sengupta, Sampurna Ghosh

Academy of Scientific and Innovative Research (AcSIR), Campus at CSIR - Central Road Research Institute, Mathura Road, New Delhi - 110 025, kunalray@gmail.com (KR); University of Calcutta, Department of Genetics, 35, Ballygunge Circular Road, Kolkata - 700 019, sengupta.mainak@gmail.com; sampurna_ghosh@yahoo.in (MS, SG) India.

Published in Atlas Database: April 2016

Online updated version : <http://AtlasGeneticsOncology.org/Genes/OCA2ID45789ch15q12.html>

Printable original version : <http://documents.irevues.inist.fr/bitstream/handle/2042/66945/04-2016-OCA2ID45789ch15q12.pdf>

DOI: 10.4267/2042/66945

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 2016 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

OCA2 gene (OCA2), having a chromosomal location of 15q12-q13, encodes an integral membrane transporter protein playing a role in regulating the pH of melanosomes.

OCA2 is hypothesized to be involved in the transport of tyrosine, the precursor to melanin synthesis, within the melanocyte. Defects in this gene are the cause of oculocutaneous albinism type II; OCA II.

Keywords: OCA2, albinism, OCA II

Identity

Other names: SHEP1, EYCL2, EYCL3, BOCA, BEY1, BEY2, EYCL, HCL3, PED, BEY, D15S12, PEDH

HGNC (Hugo): OCA2

Location: 15q12

DNA/RNA

Description


In Chromosome: 15, the 344,486 bases long gene starts from 27,754,873 bp from pter and ends at 28,099,358 bp from pter; Orientation: Minus strand. It contains 24 exons and spans ~344.5 kb of the genome.

Transcription

The gene encodes a 3154 bp transcript.

Alternative splicing results in at least two transcript variants.

Variant 2 lacks an alternate in-frame exon in the central coding region, compared to variant 1, resulting in an isoform that is shorter than isoform 1.



Cytogenetic band showing OCA2 locus (Ref: <http://www.genecards.org/cgi-bin/carddisp.pl?gene=OCA2&keywords=OCA2>)

Protein

Description

The gene encodes a protein containing 838 amino acids with molecular mass of 92850 Da. The OCA2 is thought to be a melanosomal multipass integral membrane protein (with 12 predicted transmembrane domains). OCA2 is characterized by the presence of a conserved consensus acidic dileucine-based sorting motif within the cytoplasmic N-terminal region. A second dileucine signal within this region confers steady-state lysosomal localization in melanocytes. It belongs to the CitM (TC 2.A.11) transporter family.

Expression

Due to its localization in the melanosomal membrane, OCA2 is thought to be expressed in the melanocytes.

Localisation

OCA2 is hypothesized to be present in the melanosomal membrane of the melanocytes.

Function

The precise function of OCA2 has not been elucidated till date. However, the potential functions include: a) normal biogenesis of melanosomes, b) for normal processing and transport of tyrosinase and other melanosomal proteins, and c) maintenance of an acidic pH in melanosomes.

Homology

Its sequence predicts that OCA2 has a homology to a superfamily of permeases. It has been proposed that OCA2 also regulates the post-translational processing of tyrosinase, which catalyzes the rate limiting steps in melanin biosynthesis and is a major determinant of brown and/or blue eye color.

Mutations

Germinal

Mutations in OCA2 are responsible for albinism known also as OCA2. A few OCA2 mutations have been associated also with autosomal recessive ocular albinism. While the degree of cutaneous pigment and iris color may vary, the newborn with OCA2 nearly always have pigmented hair. Nevi and freckles are common. Visual acuity is better than in OCA1 and reaches 3/10. Africans with OCA2 appear with light brown hair and skin, and gray irises. Eighty six mutations in OCA2 have been reported in Albinism Database (<http://www.ifpcs.org/albinism/oca2mut.html>). It is to be noted that Albinism Database has been updated till 2009.

Somatic

Somatic mutations in OCA2 have been identified in cancers (<http://www.cancerindex.org/geneweb/OCA2.htm>,

<http://cancer.sanger.ac.uk/cosmic/search?q=OCA2>), but no causality have been reported.

Implicated in

Melanoma

Note

In 2005, Jannot et al reported, based on allelic distribution between cases and controls, that malignant melanoma and OCA2 are associated (p value=0.030 after correction for multiple testing). In 2010 Duffy et al claimed the OCA2 variant Arg419Gln (rs1800407) to be a significant risk factor for cutaneous malignant melanoma based on a genome wide association study (GWAS). In 2011, another GWAS identified a locus at chromosome 15q13.1 (HERC2/OCA2 region) in a discovery cohort of 1804 melanoma cases and 1026 controls, to be associated with melanoma (Amos et al., 2011). Co-segregation analysis in an OCA2 affected pedigree containing individuals diagnosed with both cutaneous and iris melanoma, revealed that OCA2 variants could act as contributors towards melanoma predisposition (Hawkes et al., 2013).

The OCA2 Arg419Gln SNP has also been found to be associated with basal cell carcinoma of skin (OR, 1.50; 95% CI, 1.06-2.13) (Nan et al., 2009)

References

- Amos CI, Wang LE, Lee JE, Gershenwald JE, Chen WV, Fang S, Kosoy R, Zhang M, Qureshi AA, Vattathil S, Schacherer CW, Gardner JM, Wang Y, Bishop DT, Barrett JH, MacGregor S, Hayward NK, Martin NG, Duffy DL, Mann GJ, Cust A, Hopper J, Brown KM, Grimm EA, Xu Y, Han Y, Jing K, McHugh C, Laurie CC, Doheny KF, Pugh EW, Seldin MF, Han J, Wei Q. Genome-wide association study identifies novel loci predisposing to cutaneous melanoma. *Hum Mol Genet.* 2011 Dec 15;20(24):5012-23
- Duffy DL, Zhao ZZ, Sturm RA, Hayward NK, Martin NG, Montgomery GW. Multiple pigmentation gene polymorphisms account for a substantial proportion of risk of cutaneous malignant melanoma. *J Invest Dermatol.* 2010 Feb;130(2):520-8
- Hawkes JE, Cassidy PB, Manga P, Boissy RE, Goldgar D, Cannon-Albright L, Florell SR, Leachman SA. Report of a novel OCA2 gene mutation and an investigation of OCA2 variants on melanoma risk in a familial melanoma pedigree. *J Dermatol Sci.* 2013 Jan;69(1):30-7
- Jannot AS, Meziani R, Bertrand G, Gérard B, Descamps V, Archimbaud A, Picard C, Ollivaud L, Basset-Seguin N, Kerob D, Lanternier G, Lebbe C, Saiag P, Crickx B, Clerget-Darpoux F, Grandchamp B, Soufir N, Melan-Cohort. Allele variations in the OCA2 gene (pink-eyed-dilution locus) are associated with genetic susceptibility to melanoma *Eur J Hum Genet* 2005 Aug;13(8):913-20
- Kim HK, Kim YS, Lee SH, Lee HH. Impact of a Delayed Laparoscopic Appendectomy on the Risk of Complications in Acute Appendicitis: A Retrospective Study of 4,065 Patients *Dig Surg* 2016 Jul 6;34(1):25-29
- Kunnus K, Zhang W, Delcey MG, Pinjari RV, Miedema PS, Schreck S, Quevedo W, Schroeder H, Foehlich A, Gaffney KJ, Lundberg M, Odelius M, Wernet P. Viewing the Valence

Electronic Structure of Ferric and Ferrous Hexacyanide in Solution From the Fe and Cyanide Perspectives J Phys Chem B 2016 Jul 5

Nan H, Kraft P, Hunter DJ, Han J. Genetic variants in pigmentation genes, pigmentary phenotypes, and risk of skin cancer in Caucasians Int J Cancer 2009 Aug 15;125(4):909-17

This article should be referenced as such:

Ray K, Sengupta M, Ghosh S. OCA2 (OCA2 melanosomal transmembrane protein). Atlas Genet Cytogenet Oncol Haematol. 2016; 20(12):592-594.
