

Gene Section

Review

FANCL (FA complementation group L)

Sylvie van Twest, Andrew Deans

St Vincent's Institute of Medical Research, 9 Princes St, Fitzroy VIC 3065; svantwest@svi.edu.au
adeans@svi.edu.au

Published in Atlas Database: June 2019

Online updated version : <http://AtlasGeneticsOncology.org/Genes/FANCLID40529ch2p16.html>

Printable original version : <http://documents.irevues.inist.fr/bitstream/handle/2042/70722/06-2019-FANCLID40529ch2p16.pdf>

DOI: 10.4267/2042/70722

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2020 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

FANCL is the catalytically active component of the Fanconi anemia (FA) DNA repair pathway that maintains genomic stability by recognizing and repairing interstand cross links (ICL), and DNA damage incurred during replication.

The FA pathway is comprised of 22 genes, biallelic mutations in any one of these genes causes Fanconi anemia, a cancer pre-disposition syndrome characterized by chromosomal instability and hypersensitivity to DNA crosslinking agents, such as those used in chemotherapy like mitomycin C (MMC) (Niraj, Färkkilä et al., 2019).

FANCL acts within the 9 protein FA "core complex" (FANCA, FANCG, FAAP20 (AG20), FANCC, FANCE, FANCF (CEF), FANCB, FANCL, FAAP100 (BL100) that forms in response to DNA damage. Together with E2 conjugating enzyme Ube2t (FANCT), the E3 RING ligase FANCL monoubiquitinates FANCD2 and FANCI (ID2), this

signals downstream repair processes, and is defective in 95% of all FA patients (Inc, 2014).

Keywords

FANCL, RING E3 ligase, Fanconi Anemia, ubiquitination, cancer pre-disposition

Identity

Other names: PHF9, FAAP43, POG

HGNC (Hugo): FANCL

Location

FANCL is located on 2p16.1 which is the short arm (p) of chromosome 2 at position 16.1 between base pairs 58,159,243 to 58,241,681.

DNA/RNA

FANCL has 2 described isoforms produced by alternative splicing. Isoform one (Q9NW38-1). Is known as the canonical isoform, while isoform 2 (Q9NW38-2) differs from the canonical sequence at 178-178: T--> TPQVNS.




Figure 1: Genomic context of FANCL on chromosome 2 (Adapted from NCBI).



Figure 2: Exons in FANCL gene. Colour coded to indicate ELF, DRWD, and RING finger domains. Adapted from Chandrasekharappa et al. 2013.

Protein

Description

The FANCL gene encodes FANCL protein comprised of 375 amino acids with a molecular mass of 42905 Da. FANCL is comprised of 3 domains, an N-terminal E2-like fold (ELF), a novel double-RWD (DRWD) and C-terminal RING domain (Hodson, Purkiss et al., 2014).

Expression

From total RNA sequencing, FANCL was found to be expressed in adrenal gland (RPKM 2.1), prostate (RPKM 2.34), thymus (RPKM 2.1), and thyroid (RPKM 2.2) (Bioproject PRJNA280600 (PMID 25970244)).

In another RNA sequencing project on 27 different tissues from 95 human individuals, FANCL was highly expressed in adrenal gland (RPKM 16.8),

endometrium (RPKM 10.6), lymph nodes (RPKM 8.5), ovary (RPKM 9.2), prostate (RPKM 8.5), and testis (RPKM 12) (Bioproject PRJEB4337, PMID 24309898).

Function

FANCL the catalytically active part of the 9 protein Fanconi anemia (FA) core complex comprised of FANCB, FAAP100, FANCA, FANCG, FAAP20, FANCC, FANCE and FANCF that forms in response to DNA damage incurred during DNA replication in S-phase, or to detection of interstrand cross links (ICL) (Ceccaldi, Sarangi et al., 2016). FANCL is an E3 ubiquitin ligase that specifically monoubiquitinates FANCD2 (at lysine 561) and FANCI (at lysine 523) (ID2; Note the FANCD2-FANCI heterodimer "ID2" must not be confused with the gene ID2) in the presence of UBE2T (FANCT) to signal downstream DNA repair proteins.




Figure 3: A) Schematic of D2 monoubiquitination by E3 RING ligase FANCL and E2 conjugating enzyme. B) Ribbon diagram of FANCL with ELF domain (brown), DRWD domain (green), and RING domain (green). C) Surface representation of protein binding domains on FANCL. The binding patch for ubiquitin (orange) is within the ELF domain, while the substrate binding domain (red) is in the DRWD domain, and the Ube2t binding domain (light purple) is in the RING domain. Figure from Specificity and disease in the ubiquitin system by Viduth K. Chaugule and Helen Walden in Biochemical Society Transactions Feb 2016, 44 (1) 212-227; DOI: 10.1042/BST20150209.




Figure 4: Schematic of Fanconi Anemia DNA damage response pathway. In response to interstrand cross links (ICL), or DNA damage from DNA replication, FANCM recruits the 9 protein core complex to DNA damage sites to monoubiquitinate FANCL D2 and I. The core complex is comprised of 3 sub-complexes AG20 (FANCA, G, FAAP20), BL100 (FANCB, L, FAAP100), and CEF (FANCC, E, F). Dashed lines indicate groupings of sub-complexes, while triple lines indicate putative direct protein interactions. Within the core complex, FANCL has a RING E3 domain with ubiquitin ligase activity, but mutation in any one of the FA genes leads to defective DNA repair. Ubiquitinated ID2 is activated, and localized to chromatin in nuclear foci to interact with downstream DNA repair proteins (FANCD1, FANCD1, FANCN) to repair DNA via homologous recombination. Once DNA repair is completed, USP1 deubiquitinates ID2 so that DNA damage response can be reinitiated. Figure adapted from <https://www.sciencedirect.com/topics/biochemistry-genetics-and-molecular-biology/fancl>

FANCL is comprised of 3 distinct functional domains: the RING domain interacts with the E2 conjugating enzyme UBE2T (FANCT), the central DRWD domain interacts with FANCD2, and the N-terminal E2-like fold domain (ELF) domain interacts with ubiquitin (Hodson et al., 2014, Miles, Frost et al., 2015).

Within the core complex, FANCL interacts as a subcomplex with FANCB and FAAP100 (Huang, Leung et al., 2014, van Twest, Murphy et al., 2017); both proteins stabilize FANCL (Rajendra, Oestergaard et al., 2014), and enhance its activity 5-fold in vitro assays (Ling, Ishiai et al., 2007).

Along with FANCA, FANCG, FANCF, FANCL was found to interact directly with hairy enhancer of split 1 (HES1), which is a part of the NOTCH1 developmental pathway involved in hematopoietic stem cell (HSC) self-renewal (Tremblay, Huang et al., 2008).

Depletion of HES1 from cells resulted in FA-like phenotype with disrupted interaction between individual core complex proteins, increased cell sensitivity to DNA crosslinking agents, and reduced MMC-induced ID2 monoubiquitination (Tremblay et al., 2008).

Finally HES1 did not interact with FA-mutated core complex proteins. HSC defects and eventual bone marrow failure in FA patients may be linked to inability of HES1 to interact with a defective core

complex (Tremblay et al., 2008, Tremblay, Huang et al., 2018).

Mutations

Germinal

FANCL-associated Fanconi anemia is inherited in an autosomal recessive manner, and accounts for 0.2% of all FA cases (Wu, Liu et al., 2017). To date, there are only 9 documented cases of FANCL-associated FA (Ali, Kirby et al., 2009, Ameziane, Sie et al., 2012, Chandrasekharappa, Lach et al., 2013, Meetei, de Winter et al., 2003, Vetro, Iacone et al., 2015, Wu et al., 2017).

Of the 5 cases with phenotypic and genotypic data, four were severe, and one was mild. Two severe cases frame shift deletions in exon 4 and 6 that truncated FANCL, resulted in postnatal mortality and presented with VACTERL association (vertebral defects, anal atresia, cardiac defects, tracheoesophageal fistula, renal malformations, and limb defects) (Vetro et al., 2015). Another case with homozygous frameshift insertion of exon 9 had a severe phenotype with esophageal atresia (Ameziane et al., 2012). Finally, a novel homozygous mutation c.822_823insCTTTCAGG (p.Asp275LeufsX13) had a typical FA presentation with progression to bone marrow failure and death at age 9 from acute myelomonocytic leukemia (AML-M4) (Wu et al.,

2017). The patient with mild FANCL-associated FA had bi-allelic mutation (Ali et al., 2009). One allele had an in-frame 3-bp deletion c.1007_1009delTAT (p.Ile336_Cys337delinsSer) in exon 12 within the PHD/RING-finger domain that resulted in loss of one amino acid, isoleucine-336, and conversion of cysteine-337 to serine null mutation (produces non-function protein). The other mutated allele had a 4-bp duplication (c.1095_1098dupAATT (p.Thr367AsnfsX13) that resulted in a frameshift just outside the RING-finger domain in exon 14. 4-bp duplication mutation is a hypomorphic mutation (produces partially functional protein) (Ali et al., 2009).

Sequencing screen of 27 FA cases with unidentified mutations uncovered 3 FANCL FA associated mutations: c.375-2033C>G (skips exons 4,6,7), c.375-2033 C>G (multiple splicing aberrations), c.1092G>A (skips exon 13), but didn't have corresponding phenotypic data (Chandrasekharappa et al., 2013). There is no phenotypic data for the first FANCL-associated FA patient that had exon 11 deletion and insertion of 177-nt sequence (Meetei et al., 2003).

Implicated in

Fanconi Anemia

Disease

Biallelic mutations in FANCL, or any of the other 21 FA pathway proteins is implicated in Fanconi Anemia (FA), a rare genetic condition that results in progressive bone marrow failure (pancytopenia), congenital malformations in 75% of patients (short stature, urogenital defects, café au lait spots, skeletal malformations), and cancer pre-disposition (primarily acute myeloid leukaemia, and certain solid tumours) (Alter, 2014).

Mutations that result in loss-of-function of both FANCL alleles may correlate with more severe phenotypes (Vetro et al., 2015). Hydrocephalus-VACTERL (vertebral, anal, cardiac, tracheo-esophageal fistula, renal, and limb anomalies) syndrome has been reported in two FANCL-linked FA patients that died shortly after birth (Vetro et al., 2015).

Prognosis: The prognosis for FA is poor as there is no cure, and the average lifespan is 20-30 years. If no congenital abnormalities are apparent at birth, patients are often diagnosed with FA when they present with aplastic anemia ages 8-10 (>700 fold risk) (Alter, 2014). Bone marrow transplants are often conducted to correct the haematological issues associated with FA, however due to faulty DNA repair FA patients retain high cancer risk particularly leukaemia, and head and neck squamous cell carcinomas (approximately 500 fold risk) (Shimamura Alter, 2010).

Diagnosis: Diagnostics for FA is done with a chromosomal breakage test; when treated with interstrand crosslinking agents such as mitomycin C (MMC) or diepoxybutane (DEB) FA cells exhibit high number chromosomal breakages, and abnormalities as compared to normal cells.

References

- Abramson DH, Frank CM, Susman M, Whalen MP, Dunkel IJ, Boyd NW 3rd. Presenting signs of retinoblastoma. *J Pediatr*. 1998 Mar;132(3 Pt 1):505-8
- Albrecht P, Ansperger-Rescher B, Schüler A, Zeschnigk M, Gallie B, Lohmann DR. Spectrum of gross deletions and insertions in the RB1 gene in patients with retinoblastoma and association with phenotypic expression. *Hum Mutat*. 2005 Nov;26(5):437-45
- Blanquet V, Turleau C, Créau-Goldberg N, Cochet C, de Grouchy J. De novo t(2;13)(p24.3;q14.2) and retinoblastoma. Mapping of two 13q14 probes by in situ hybridization. *Hum Genet*. 1987 May;76(1):102-5
- Broadus E, Topham A, Singh AD. Incidence of retinoblastoma in the USA: 1975-2004 *Br J Ophthalmol* 2009 Jan;93(1):21-3
- Caselli R, Speciale C, Pescucci C, Uliana V, Sampieri K, Bruttini M, Longo I, De Francesco S, Pramparo T, Zuffardi O, Frezzotti R, Acquaviva A, Hadjistilianou T, Renieri A, Mari F. Retinoblastoma and mental retardation microdeletion syndrome: clinical characterization and molecular dissection using array CGH *J Hum Genet* 2007;52(6):535-42
- Cavenee WK, Dryja TP, Phillips RA, Benedict WF, Godbout R, Gallie BL, Murphree AL, Strong LC, White RL. Expression of recessive alleles by chromosomal mechanisms in retinoblastoma *Nature* 1983 Oct 27-Nov 2;305(5937):779-84
- Cross HE, Hansen RC, Morrow G 3rd, Davis JR. Retinoblastoma in a patient with a 13qXp translocation *Am J Ophthalmol* 1977 Oct;84(4):548-54
- Dehainault C, Garancher A, Castéra L, Cassoux N, Aerts I, Doz F, Desjardins L, Lumbroso L, Montes de Oca R, Almouzni G, Stoppa-Lyonnet D, Pouponnot C, Gauthier-Villars M, Houdayer C. The survival gene MED4 explains low penetrance retinoblastoma in patients with large RB1 deletion *Hum Mol Genet* 2014 Oct 1;23(19):5243-50
- Dimaras H, Corson TW, Cobrinik D, White A, Zhao J, Munier FL, Abramson DH, Shields CL, Chantada GL, Njuguna F, Gallie BL. Retinoblastoma *Nat Rev Dis Primers* 2015 Aug 27;1:15021
- Dries D, Baca K, Truss L, Dobin S. Interstitial deletion of 13q and a 13;X chromosome translocation results in partial trisomy 13 and bilateral retinoblastoma *Ophthalmic Genet* 2003 Sep;24(3):175-80
- Dyson NJ. RB1: a prototype tumor suppressor and an enigma *Genes Dev* 2016 Jul 1;30(13):1492-502
- Eloy P, Dehainault C, Sefta M, Aerts I, Doz F, Cassoux N, Lumbroso le Rouic L, Stoppa-Lyonnet D, Radvanyi F, Millot GA, Gauthier-Villars M, Houdayer C. A Parent-of-Origin Effect Impacts the Phenotype in Low Penetrance Retinoblastoma Families Segregating the c 1981C>T/p

Arg661Trp Mutation of RB1

- Friend SH, Bernards R, Rogelj S, Weinberg RA, Rapaport JM, Albert DM, Dryja TP. A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma *Nature* 1986 Oct 16-22;323(6089):643-6
- Greger V, Debus N, Lohmann D, Höpping W, Passarge E, Horsthemke B. Frequency and parental origin of hypermethylated RB1 alleles in retinoblastoma *Hum Genet* 1994 Nov;94(5):491-6
- Hagstrom SA, Dryja TP. Mitotic recombination map of 13cen-13q14 derived from an investigation of loss of heterozygosity in retinoblastomas *Proc Natl Acad Sci U S A* 1999 Mar 16;96(6):2952-7
- Huddleston S, McNall-Knapp RY, Siatkowski M, Odom C, Brennan R, Wilson MW. A novel translocation t(11;13)(q21;q14 2) in a child with suprasellar primitive neuroectodermal tumor and retinoblastoma *Ophthalmic Genet*
- Imperatore V, Pinto AM, Gelli E, Trevisson E, Morbidoni V, Frullanti E, Hadjistilianou T, De Francesco S, Toti P, Gusson E, Roversi G, Accogli A, Capra V, Mencarelli MA, Renieri A, Ariani F. Parent-of-origin effect of hypomorphic pathogenic variants and somatic mosaicism impact on phenotypic expression of retinoblastoma *Eur J Hum Genet* 2018 Jul;26(7):1026-1037
- Inc F ni. Fanconi Anemia: Guidelines for Diagnosis and Management. Fanconi Anemia Research Fund (2014)
- Kajii T, Tsukahara M, Fukushima Y, Hata A, Matsuo K, Kuroki Y. Translocation (X;13)(p11 21;q12 3) in a girl with incontinentia pigmenti and bilateral retinoblastoma
- Keith CG, Webb GC. Retinoblastoma and retinoma occurring in a child with a translocation and deletion of the long arm of chromosome 13 *Arch Ophthalmol* 1985 Jul;103(7):941-4
- Kloss K, Währisch P, Greger V, Messmer E, Fritze H, Höpping W, Passarge E, Horsthemke B. Characterization of deletions at the retinoblastoma locus in patients with bilateral retinoblastoma *Am J Med Genet* 1991 May 1;39(2):196-200
- Klutzm M, Horsthemke B, Lohmann DR. RB1 gene mutations in peripheral blood DNA of patients with isolated unilateral retinoblastoma *Am J Hum Genet* 1999 Feb;64(2):667-8
- Knudson AG Jr. Mutation and cancer: statistical study of retinoblastoma *Proc Natl Acad Sci U S A* 1971 Apr;68(4):820-3
- Laquis SJ, Rodriguez-Galindo C, Wilson MW, Fleming JC, Haik BG. Retinoblastoma in a patient with an X;13 translocation and facial abnormalities consistent with 13q-syndrome *Am J Ophthalmol* 2002 Feb;133(2):285-7
- Lohmann DR, Gerick M, Brandt B, Oelschläger U, Lorenz B, Passarge E, Horsthemke B. Constitutional RB1-gene mutations in patients with isolated unilateral retinoblastoma *Am J Hum Genet* 1997 Aug;61(2):282-94
- Ohtani-Fujita N, Dryja TP, Rapaport JM, Fujita T, Matsumura S, Ozasa K, Watanabe Y, Hayashi K, Maeda K, Kinoshita S, Matsumura T, Ohnishi Y, Hotta Y, Takahashi R, Kato MV, Ishizaki K, Sasaki MS, Horsthemke B, Minoda K, Sakai T. Hypermethylation in the retinoblastoma gene is associated with unilateral, sporadic retinoblastoma *Cancer Genet Cytogenet* 1997 Oct 1;98(1):43-9
- Peterson EC, Elhammady MS, Quintero-Wolfe S, Murray TG, Aziz-Sultan MA. Selective ophthalmic artery infusion of chemotherapy for advanced intraocular retinoblastoma: initial experience with 17 tumors *J Neurosurg* 2011 Jun;114(6):1603-8
- Triviño E, Guitart M, Egozcue J, Coll MD. Characterization by FISH of a t(5;13) in a patient with bilateral retinoblastoma *Cancer Genet Cytogenet* 1997 Jul 1;96(1):23-5
- Venturi C, Bracco S, Cerase A, Cioni S, Galluzzi P, Gennari P, Vallone IM, Tinturini R, Vittori C, De Francesco S, Caini M, D'Ambrosio A, Toti P, Renieri A, Hadjistilianou T. Superselective ophthalmic artery infusion of melphalan for intraocular retinoblastoma: preliminary results from 140 treatments *Acta Ophthalmol* 2013 Jun;91(4):335-42
- Vogel F. Genetics of retinoblastoma *Hum Genet* 1979 Nov 1;52(1):1-54
- Zhu X, Dunn JM, Goddard AD, Squire JA, Becker A, Phillips RA, Gallie BL. Mechanisms of loss of heterozygosity in retinoblastoma *Cytogenet Cell Genet* 1992;59(4):248-52

This article should be referenced as such:

van Twest S, Deans A. FANCL (FA complementation group L). *Atlas Genet Cytogenet Oncol Haematol.* 2020; 24(5):189-193.
